



## ACETAMINOPHEN

Brand Names: Tylenol

Weight (lbs)	Infant or Children's Liquid Suspension (160 mg/5 mL) Use dosing cup or oral syringe provided	Chewable Tablets (80 mg)	Chewable Tablets (160 mg)	Tablets (325 mg)
6-11 lbs	1.25 mL	—	—	—
12-17 lbs	2.5 mL	—	—	—
18-23 lbs	3.75 mL	1.5 tablets	—	—
24-35 lbs	5 mL	2 tablets	1 tablet	—
36-47 lbs	7.5 mL	3 tablets	1.5 tablets	—
48-59 lbs	10 mL	4 tablets	2 tablets	1 tablet
60-71 lbs	12.5 mL	5 tablets	2.5 tablets	1 tablet
72-95 lbs	15 mL	6 tablets	3 tablets	1.5 tablets
96+ lbs	20mL	8 tablets	4 tablets	2 tablets

### INSTRUCTIONS:

- Always check & read the label on the medication bottle.
- May give every 4-6 hours as needed. **Do not give more than 5 doses** in 24 hours.
- Do not use with other products/medications containing acetaminophen.

### AGE LIMITS:

- Do not give acetaminophen to a baby under 3 months of age. Infants under 3 months old with a fever over 100.4 degrees (38°C) rectally should be seen by a healthcare provider.
- Chewable tablets are not recommended for children under 2 years old.

## IBUPROFEN

Brand Names: Motrin, Advil

Weight (lbs)	Infant's Liquid Suspension (50 mg/1.25 mL) Use dropper provided	Children's Liquid Suspension (100 mg/5 mL) Use dosing cup or oral syringe provided	Junior Tablets (100 mg)	Tablets (200 mg)
6-11 lbs	Consult provider		—	—
12-17 lbs	1.25 mL	2.5 mL	—	—
18-23 lbs	1.875 mL	3.75 mL	—	—
24-35 lbs	—	5 mL	1 tablet	—
36-47 lbs	—	7.5 mL	1.5 tablets	—
48-59 lbs	—	10 mL	2 tablets	1 tablet
60-71 lbs	—	12.5 mL	2.5 tablets	1 tablet
72-95 lbs	—	15 mL	3 tablets	1.5 tablets
96+ lbs	—	20 mL	4 tablets	2 tablets

### INSTRUCTIONS:

- Always check & read the label on the medication bottle.
- May give every 6-8 hours as needed. **Do not give more than 4 doses** in 24 hours.

### AGE LIMITS:

- Do not give ibuprofen to a baby under 6 months of age.
- Chewable tablets are not recommended for children under 2 years old.



UTAH POISON CONTROL CENTER



## ACETAMINOFÉN

Nombres de marca: Tylenol, Panadol (paracetamol)

Peso (libras)	Suspensión líquida para bebés o niños (160 mg / 5 mL) Use el vaso dosificador o la jeringa oral provista	Tabletas masticables (80 mg)	Tabletas masticables (160 mg)	Tabletas (325 mg)
6-11 libras	1.25 mL	—	—	—
12-17 libras	2.5 mL	—	—	—
18-23 libras	3.75 mL	1.5 tabletas	—	—
24-35 libras	5 mL	2 tabletas	1 tableta	—
36-47 libras	7.5 mL	3 tabletas	1.5 tabletas	—
48-59 libras	10 mL	4 tabletas	2 tabletas	1 tableta
60-71 libras	12.5 mL	5 tabletas	2.5 tabletas	1 tableta
72-95 libras	15 mL	6 tabletas	3 tabletas	1.5 tabletas
96+ libras	20mL	8 tabletas	4 tabletas	2 tabletas

### INSTRUCCIONES:

- Siempre revise y lea la etiqueta en el frasco del medicamento.
- Puede administrar cada 4-6 horas según sea necesario. **No administre más de 5 dosis** en 24 horas.
- No lo use con otros productos/medicamentos que contengan acetaminofén.

### LÍMITES DE EDAD:

- No le dé acetaminofén a un bebé menor de 3 meses de edad. Los bebés menores de 3 meses con fiebre rectal superior a 100.4 grados (38°C) deben ser atendidos por un proveedor de atención médica.
- Las tabletas masticables no se recomiendan para niños menores de 2 años.

## IBUPROFENO

Nombres de marca: Motrin, Advil

Weight (libras)	Suspensión líquida para bebés (50 mg/1.25 mL) Utilice el cuentagotas suministrado	Suspensión líquida para niños (100 mg/5 mL) Use el vaso dosificador o la jeringa oral provista	Tabletas Junior (100 mg)	Tabletas (200 mg)
6-11 libras	Consultar proveedor		—	—
12-17 libras	1.25 mL	2.5 mL	—	—
18-23 libras	1.875 mL	3.75 mL	—	—
24-35 libras	—	5 mL	1 tableta	—
36-47 libras	—	7.5 mL	1.5 tabletas	—
48-59 libras	—	10 mL	2 tabletas	1 tableta
60-71 libras	—	12.5 mL	2.5 tabletas	1 tableta
72-95 libras	—	15 mL	3 tabletas	1.5 tabletas
96+ libras	—	20 mL	4 tabletas	2 tabletas

### INSTRUCCIONES:

- Siempre revise y lea la etiqueta en el frasco del medicamento.
- Puede administrar cada 6-8 horas según sea necesario. **No administre más de 4 dosis** en 24 horas.

### LÍMITES DE EDAD:

- Siempre revise y lea la etiqueta en el frasco del medicamento.
- Puede administrar cada 6-8 horas según sea necesario. **No administre más de 4 dosis** en 24 horas.



UTAH POISON  
CONTROL CENTER